

Androgen Receptor (9Q12) Rabbit Monoclonal Antibody

Catalog# UZ-MA00298

Product information

产品货号	UZ-MA00298
别名	KD; AIS; AR8; TFM; DHTR; SBMA; HYSPI; NR3C4; SMAX1; HUMARA
产品名称	Androgen Receptor (9Q12) Rabbit Monoclonal Antibody
类别	抗体产品
推荐应用	WB
反应种属	Human,Mouse,Rat
存储缓冲液	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
免疫原	A synthetic peptide of human Androgen Receptor
稀释度	WB 1:1000-1:5000
参考分子量	110kDa
预测分子量	99kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	IgG
背景介绍	The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and an androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (SBMA, also known as Kennedy's disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2017]
纯化	Affinity Purification
Uniprot ID	P10275
Clonality	Monoclonal