

53BP1 Rabbit Polyclonal Antibody

Catalog# UZ-PA00081

Product information

产品货号	UZ-PA00081
别名	TP53BP1; Tumor suppressor p53-binding protein 1; 53BP1; p53-binding protein 1; p53BP1
产品名称	53BP1 Rabbit Polyclonal Antibody
类别	抗体产品
基因名称	TP53BP1
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Rat,Mouse
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N
Human Gene ID	7158
Human Swissprot No.	Q12888
免疫原	The antiserum was produced against synthesized peptide derived from human TP53B P1. AA range:1744-1793
稀释度	WB 1:500-1:2000, IHC-P 1:100-1:300, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:10000 .Not yet tested in other applications.
参考分子量	214kDa
运输及保存条件	-20°C/1 year
宿主	Rabbit
同种型	Rabbit,IgG
背景介绍	function:May have a role in checkpoint signaling during mitosis (By similarity). Enhances TP53-mediated transcriptional activation. Plays a role in the response to DNA damage.,PTM:Asymmetrically dimethylated on Arg residues by PRMT1. Methylation is required for DNA binding.,PTM:Phosphorylated at basal level in the absence of DNA damage. Hyper-phosphorylated in an ATM-dependent manner in response to DNA damage induced by ionizing radiation. Hyper-phosphorylated in an ATR-dependent manner in response to DNA damage induced by UV irradiation.,similarity:Contains 2 BRCT domains.,subcellular location:Associated with kinetochores. Both nuclear and cytoplasmic in some cells. Recruited to sites of DNA damage, such as double strand breaks. Methylation of histone H4 at 'Lys-20' is required for efficient localization to double strand breaks.,subunit:Interacts with IFI202A (By similarity). Binds to the central domain of TP53/p53. May form homo-oligomers. Interacts with DCLRE1C. Interacts with histone H2AFX and this requires phosphorylation of H2AFX on 'Ser-139'. Interacts with histone H4 that has been dimethylated at 'Lys-20'. Has low affinity for histone H4 containing monomethylated 'Lys-20'. Does not bind histone H4 containing unmethylated or trimethylated 'Lys-20'. Has low affinity for histone H3 that has been dimethylated on 'Lys-79'. Has very low affinity for histone H3 that has been monomethylated on 'Lys-79' (in vitro). Does not bind unmethylated histone H3.,
细胞定位	Nucleus . Chromosome . Chromosome, centromere, kinetochore . Localizes to the nucleus in absence of DNA damage (PubMed:28241136). Following DNA damage, recruited to sites of DNA damage, such as double strand breaks (DSBs): recognizes and binds histone H2A monoubiquitinated at 'Lys-15' (H2AK15Ub) and histone H4 dimethylated at 'Lys-20' (H4K20me2), two histone marks that are present at DSBs sites (PubMed:23333306, PubMed:23760478, PubMed:24703952, PubMed:28241136, PubMed:17190600). Associated with kinetochores during mitosis (By similarity). .

53BP1 Rabbit Polyclonal Antibody

Catalog# UZ-PA00081 (cont.)

Product information

纯化	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal