

Chk2 (17Y16) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00038

Product information

产品货号	HZ-YMA00038
别名	CHEK2;CDS1;CHK2;RAD53;Serine/threonine-protein kinase Chk2;CHK2 checkpoint homolog;Cds1 homolog;Hucds1;hCds1;Checkpoint kinase 2
产品名称	Chk2 (17Y16) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	CHEK2
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	11200
Human Swissprot No.	O96017
稀释度	IHC-P 1:1000-1:4000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:500-1:20000, IP 1:50-1:200
参考分子量	61kDa
预测分子量	61kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations
纯化	Protein A
Clonality	Monoclonal