

eIF2 α (9E12) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00043

Product information

产品货号	HZ-YMA00043
别名	EIF2S1;EIF2A;Eukaryotic translation initiation factor 2 subunit 1;Eukaryotic translation initiation factor 2 subunit alpha;eIF-2-alpha;eIF-2A;eIF-2alpha
产品名称	eIF2 α (9E12) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	EIF2S1
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	1965
Human Swissprot No.	P05198
稀释度	IHC-P 1:1000-1:5000, WB 1:2000-1:10000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	36kDa
预测分子量	36kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010].
纯化	Protein A
Clonality	Monoclonal