

Bag3 (15W17) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00089

Product information

产品货号	HZ-YMA00089
别名	BAG3;BIS;BAG family molecular chaperone regulator 3;BAG-3;Bcl-2-associated athanogene 3;Bcl-2-binding protein Bis;Docking protein CAIR-1
产品名称	Bag3 (15W17) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	BAG3
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	9531
Human Swissprot No.	O95817
稀释度	IHC-P 1:200-1:1000, WB 1:1000-1:5000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:500-1:20000
参考分子量	80kDa
预测分子量	62kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008],
纯化	Protein A
Clonality	Monoclonal