

RPA32/RPA2 (5C3) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00114

Product information

产品货号	HZ-YMA00114
别名	RPA2;REPA2;RPA32;RPA34;Replication protein A 32 kDa subunit;RP-A p32;Replication factor A protein 2;RF-A protein 2;Replication protein A 34 kDa subunit;RP-A p34
产品名称	RPA32/RPA2 (5C3) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	RPA2
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	6118
Human Swissprot No.	P15927
稀释度	IHC-P 1:1000-1:4000, WB 1:1000-1:5000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000, IP 1:50-1:200
参考分子量	29kDa
预测分子量	29kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	function:Required for DNA recombination, repair and replication. The activity of RP-A is mediated by single-stranded DNA binding and protein interactions.,PTM:Phosphorylated in a cell-cycle-dependent manner (from the S phase until mitosis). Phosphorylated by ATR upon DNA damage, which promotes its translocation to nuclear foci. Can be phosphorylated in vitro by PRKDC/DNA-PK in the presence of Ku and DNA, and by CDC2.,subcellular location:Also present in PML nuclear bodies. Redistributes to discrete nuclear foci upon DNA damage.,subunit:Heterotrimer of 70, 32 and 14 kDa chains. The DNA-binding activity may reside exclusively on the 70 kDa subunit. Binds to SERTAD3/RBT1. Interacts with TIPIN.,
纯化	Protein A
Clonality	Monoclonal