

Glucocorticoid Receptor (14O8) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00165

Product information

产品货号	HZ-YMA00165
别名	GCCR;GCR;GCR_HUMAN;GCRST;glucocorticoid nuclear receptor variant 1;Glucocorticoid receptor;GR;GRL;Grl1;nr3c1;Nuclear receptor subfamily 3 group C member 1;nuclear receptor subfamily 3, group C, member 1 (glucocorticoid receptor);Grl;Grl1
产品名称	Glucocorticoid Receptor (14O8) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	Nr3c1
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	2908
Human Swissprot No.	P04150
稀释度	IHC-P 1:1000-1:4000, WB 1:1000-1:5000, IF-P/IF-F/IF-ICC 1:200-1:1000, ELISA 1:5000-1:20000
参考分子量	94kDa
预测分子量	85kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175). [provided by RefSeq, Feb 2011]
纯化	Protein A
Clonality	Monoclonal