

Collagen IV (5S16) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00271

Product information

产品货号	HZ-YMA00271
别名	Arresten;BSVD;CO4A1_HUMAN;COL4A1;COL4A1 NC1 domain;COL4A2;COL4A3;COL4A4;COL4A5;collagen alpha-1(IV) chain;Collagen IV Alpha 1 Polypeptide;Collagen IV Alpha 2 Polypeptide;Collagen Of Basement Membrane Alpha 1 Chain;Collagen Of Basement Membrane Alpha 2 Chain;Collagen Type IV Alpha 1;collagen type IV alpha 1 chain;Collagen Type IV Alpha 2;Collagen Type IV Alpha 3;Collagen Type IV Alpha 4;Collagen Type IV Alpha 5;RatOR
产品名称	Collagen IV (5S16) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	COL4A1
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,ELISA
反应种属	Human
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	1282
Human Swissprot No.	P02462
稀释度	IHC-P 1:200-1000, WB 1:1000-5000, IF-P/IF-F/IF-ICC 1:200-1000, ELISA 1:5000-20000
参考分子量	200kDa
预测分子量	160kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	This gene encodes a type IV collagen alpha protein. Type IV collagen proteins are integral components of basement membranes. This gene shares a bidirectional promoter with a paralogous gene on the opposite strand. The protein consists of an amino-terminal 7S domain, a triple-helix forming collagenous domain, and a carboxy-terminal non-collagenous domain. It functions as part of a heterotrimer and interacts with other extracellular matrix components such as perlecan, proteoglycans, and laminins. In addition, proteolytic cleavage of the non-collagenous carboxy-terminal domain results in a biologically active fragment known as arresten, which has anti-angiogenic and tumor suppressor properties. Mutations in this gene cause porencephaly, cerebrovascular disease, and renal and muscular defects. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014],
纯化	Protein A
Clonality	Monoclonal