

Histone H4 (9P14) Rabbit Monoclonal Antibody

Catalog# HZ-YMA00274

Product information

产品货号	HZ-YMA00274
别名	HIST1H4A;H4/A;H4FA;HIST1H4B;H4/I;H4FI;HIST1H4C;H4/G;H4FG;HIST1H4D;H4/B;H4FB;HIST1H4E;H4/J;H4FJ;HIST1H4F;H4/C;H4FC;HIST1H4H;H4/H;H4FH;HIST1H4I;H4/M;H4FM;HIST1H4J;H4/E;H4FE;HIST1H4K;H4/D;H4FD;HIST1H4L;H4/K;H4FK;HIST1H4B;HIST1H4C;HIST1H4D;HIST1H4E;HIST1H4F;HIST1H4H;HIST1H4I;HIST1H4J;HIST1H4K;HIST1H4L;HIST2H4A;HIST2H4B;HIST4H4;
产品名称	Histone H4 (9P14) Rabbit Monoclonal Antibody
类别	抗体产品
基因名称	HIST1H4A
推荐应用	WB,IHC-P,IF-P,IF-F,IF-ICC,IP,ELISA,CHIP
反应种属	Human,Mouse,Rat
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Human Gene ID	121504/554313/8294/8359/8360/8361/8362/8363/8364/8365/8366/8367/8368/8370
Human Swissprot No.	P62805
稀释度	IHC-P 1:200-1000, WB 1:1000-5000, IF-P/IF-F/IF-ICC 1:200-1000, ELISA 1:5000-20000, IP 1:50-200, CHIP 1:50-1:100
参考分子量	11kDa
预测分子量	11kDa
运输及保存条件	Biological ice bag transportation. Store at -20°C for at least 12 months(Do not lower than -25°C). Avoid freeze/thaw cycles
宿主	Rabbit
同种型	Monoclonal, rabbit, IgG, Kappa
背景介绍	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015],
纯化	Protein A
Clonality	Monoclonal